Submission form

Client number / Barcode

LABOKL	.IN
LABOR FÜR KLINISCHE DIAGNOSTIK GMBH	& CO. KG

Equine Reproduction and Trade

PB 1810 DE-97668 Bad Kissingen Tel +49 971 72020 Fax +49 971 68546

linic Address:	Sample	Owner's addı	ess:	
Practice stamp or block letters)	☐ Whole blood	Name:		Your personal data will be used process your order according to
	□ Serum □ Plasma	-		terms for the use of data.
	□ Blood smear	First name:		You can find these terms as we information on your rights at
	SpermSwab with medium	Street:		http://laboklin.com/dataprotection
	 Swab without medium 	m Postal codo/si	tv.	
	Swab guttural pouchAspirate	FOSIAI COUE/CI	ty:	
	•	Fax / Email:		
AT No:		Tel.:		
ax / Email:				
ate and signature practice:				Signature owner
				<u> </u>
Horse Name:				
ex: F M M.N	I. Date of birth:		Date and time of sampling:	1
ample quality: Haemolytic	□ Icteric □	Lipaemic — Cer	trifuged — Cooled / Froze	en
fe Number (UELN):	C	hip-/Tattoo No.:		
eport: English	German			
Export certification	ate (27.50 €) □ I do r	ot require a certificate		email):
eason for Examination:				
Export to:		Date of e	xport:	
→ Export of semen to:				
Export of semen to:				
Export of semen to: Serology and PCR Testin		Cor	ntagious Equine Metritis (•
Serology and PCR Testin	ng	Col Send note:	swabs in AMIES transport medium with If you request both PCR and culture, one	charcoal. Cooled transport is required. P
	ng	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed .	charcoal. Cooled transport is required. Pe swab for each must be sent and clearly
Serology and PCR Testin	ng s * (c-ELISA) S/1ml	Col Send note:	swabs in AMIES transport medium with If you request both PCR and culture, one ed .	charcoal. Cooled transport is required. P
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (Th	ng s * (c-ELISA) ^{S/1ml} h.equi + B.caballi) _{EP,HP,S/0,5ml}	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed .	charcoal. Cooled transport is required. Pe swab for each must be sent and clearly : Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 C.F. ISA	ng s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed .	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (Th	ng s * (c-ELISA) ^{S/1ml} h.equi + B.caballi) _{EP,HP,S/0,5ml}	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed . Sampling location	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (Th 576 IFAT 2576 C-ELISA 1576 CFT *	ng s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml	Col Send note: label	swabs in AMIES transport medium with if you request both PCR and culture, one of . re Sampling location CEM (culture) (per swab)	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 C.F. ISA	s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed . re Sampling location CEM (culture) (per swab) 7 days	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 Left) C-ELISA 1576 CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum)	ng s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml EP,HP,S/0,5ml	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed . re Sampling location CEM (culture) (per swab) 7 days 14 days	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA	ng s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml EP,HP,S/0,5ml	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed . re Sampling location CEM (culture) (per swab) 7 days	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 Left) C-ELISA 1576 CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum)	s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml S/1ml	Col Send note: label	swabs in AMIES transport medium with If you request both PCR and culture, one ed . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito CEM PCR Profile Mare 2	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 C-ELISA 1576 CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA 1574 EVA (VNT) Latest vaccination again	s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml S/1ml S/0,5ml	Send note: labell Ma	swabs in AMIES transport medium with If you request both PCR and culture, one ed . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 CFT* 2050 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA 1331 EVA (VNT) Latest vaccination against EVA (PCR)	s * (c-ELISA)	Coi Send note: label Ma	swabs in AMIES transport medium with If you request both PCR and culture, one ed . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito CEM PCR Profile Mare 2	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The Figure 1975 CFT 1975	S * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml S/1ml S/0,5ml ainst EVA: A,Sp,EB	Coi Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito CEM PCR Prossa clitoridis, Sinus clitoridis, Sinus clitoridis	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 C-ELISA 1576 C-ELISA 1576 C-FT * 2050 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA 1311 EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 13130 Coggins Test	s * (c-ELISA)	Coi Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito CEM PCR Prossa clitoridis, Sinus clitoridis, Sinus clitoridis	charcoal. Cooled transport is required. Pe swab for each must be sent and clearly : Fossa clitoridis Sinus clitoridis medialis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The First CFT & C-ELISA CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 2030 Coggins Test 2646 C-ELISA	s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml S/1ml S/0,5ml ainst EVA: A,Sp,EB Virus / EIA S/0,5ml	Coi Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clitor CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clitor Sampling location: Sampling location: CEM (culture)	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 C-ELISA 1576 CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 1313 Coggins Test 1246 c-ELISA 2061 Glanders * (CFT)	s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml S/1ml S/0,5ml ainst EVA: A,Sp,EB Virus / EIA S/0,5ml	Coi Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clito CEM CEM CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clito CEM CEM CEM CEM CEM CEM CEM CEM (culture) (per swab)	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce Semen/Pre-ejaculatory fluid
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The First CFT & C-ELISA CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 2030 Coggins Test 2646 C-ELISA	s * (c-ELISA) S/1ml h.equi + B.caballi) EP,HP,S/0,5ml EP,HP,S/0,5ml S/1ml S/0,5ml ainst EVA: A,Sp,EB Virus / EIA S/0,5ml	Coi Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clitor CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clitor Sampling location: Sampling location: CEM (culture)	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce Semen/Pre-ejaculatory fluid
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The 1576 LFAT 1576 C-ELISA 1576 CFT * 2060 Dourine * (CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 1313 Coggins Test 1246 c-ELISA 2061 Glanders * (CFT)	s * (c-ELISA)	Coi Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one ed. re Sampling location CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clito CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clito CEM Culture) (per swab) 7 days 14 days CEM PCR Profile Stallion 1	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce Semen/Pre-ejaculatory fluid TM
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The Testin CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 3130 Coggins Test 2646 c-ELISA 2051 Glanders * (CFT) (Burkholderia mallei) 220 Salmonella Abortuseque	s * (c-ELISA)	Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clitor CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clitor CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clitor CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Stallion 1 (3 locations: Penile sheath, Urethra, Fo	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce Semen/Pre-ejaculatory fluid TM 3xTM ssa glandis)
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The Figure 1975 CFT 1975	s * (c-ELISA)	Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one ed . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clitor	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce Semen/Pre-ejaculatory fluid TM 3xTM ssa glandis) 3xTM+Sp
Serology and PCR Testin 2424 African Horse Sickness Babesia / Piroplasmosis (The Testin CFT) (Trypanosoma equiperdum) Equine Arteritis Virus / EVA (VNT) Latest vaccination against EVA (PCR) Equine Infectious Anaemia 3130 Coggins Test 2646 c-ELISA 2051 Glanders * (CFT) (Burkholderia mallei) 220 Salmonella Abortuseque	s * (c-ELISA)	Send note: label Ma	swabs in AMIES transport medium with if you request both PCR and culture, one of . CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Mare 1 (2 locations: Fossa clitoridis, Sinus clitor CEM Culture) (per swab) 3 mapling location: CEM (culture) (per swab) CEM PCR Profile Mare 2 (3 locations: Fossa clitoridis, Sinus clitor CEM (culture) (per swab) 7 days 14 days CEM PCR Profile Stallion 1 (3 locations: Penile sheath, Urethra, Forest PCR Profile Stallion 2	charcoal. Cooled transport is required. Perswab for each must be sent and clearly Fossa clitoridis Sinus clitoridis medialis Sinus clitoridis lateralis Cervix Uterus TM 2xTM ridis) 3xTM ridis, Cervix) Fossa glandis Sinus urethralis/Urethra Penile sheath/Prepuce Semen/Pre-ejaculatory fluid TM 3xTM ssa glandis) 3xTM+Sp

We are at your disposal - Feel free to call us



General Conditions: (Prices and conditions valid from July 1, 2023 to June 30, 2024)